NDC 0010-4709-01

# Sulmet®

## (sulfamethazine)

## Oblets® 2.5 g

Antibacterial

For Animal Use Only

**Net Contents:** 100 Oblets

NADA 122-271, Approved by FDA

2.5 g

**Indications:** This size Oblet is indicated for the treatment of the following diseases when caused by organisms susceptible to sulfamethazine.

**Calves:** Bacterial Pneumonia and Bovine Respiratory Disease Complex (Shipping Fever Complex) (*Pasteurella* spp.); Colibacillosis (Bacterial Scours) (*Escherichia coli*); Calf Diphtheria (*Fusobacterium necrophorum*)

**Foals:** Bacterial Pneumonia (secondary infections associated with *Pasteurella* spp.); Strangles (*Streptococcus equi*); Bacterial Enteritis (*Escherichia coli*)

**Once-a-Day Dosage by Mouth**

**First Day:** One Oblet for each 25 lb body weight. **Following Days:** One-half Oblet for each 25 lb body weight. This dosage will provide approximately 100 mg/lb body weight the first day, and 50 mg/lb body weight on each following day.

| Weight of Animal: | 25 lb | 50 lb | 75 lb | 100 lb | 125 lb |
|---|---|---|---|---|---|
| Dosage - First Day | 1 Oblet | 2 Oblets | 3 Oblets | 4 Oblets | 5 Oblets |
| Dosage - Following Days | 1/2 Oblet | 1 Oblet | 1-1/2 Oblets | 2 Oblets | 2-1/2 Oblets |

Larger size Oblets are available and are easier to use with heavier animals. Give indicated dose once daily until animal's temperature and appearance are normal. Lubricate Oblets before dosing.

**Precautions:** Sulmet sulfamethazine works fast. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Excessive dosage may cause toxic reactions. Follow above dosages carefully. Fluid intake must be adequate while treating with Sulmet. For best advice in the control and treatment of disease, consult a veterinarian. Treatment of all diseases should be instituted early. Treatment should continue 24 to 48 hours beyond the remission of disease symptoms, but not to exceed 5 consecutive days.

**Residue Warnings:** Treated cattle must not be slaughtered for at least 10 days after the last dose. Exceeding five (5) consecutive doses may cause violative tissue residue to remain beyond the withdrawal time. Do not use in calves under one (1) month of age or calves being fed an all-milk diet. Use in these classes of calves may cause violative residues to remain beyond the withdrawal time. Do not use in female dairy cattle 20 months of age or older. Use of sulfamethazine in this class of cattle may cause milk residues.

**Do not use in horses intended for human consumption.**

Restricted Drug (California) - Use Only as Directed

Store at or below 25°C (77°F).

Sulmet and Oblets are registered trademarks of Boehringer Ingelheim Vetmedica, Inc.

Manufactured for:
**Boehringer Ingelheim Vetmedica, Inc.**
St. Joseph, MO 64506 U.S.A.

12730
D6932C

Boehringer Ingelheim

Lot No.:

Exp. Date:

0 12313 47091 3