


FACT SHEET



Active Constituents: IVERMECTIN 5 mg/mL

PRODUCT DESCRIPTION

For the treatment and control of ivermectin-sensitive gastrointestinal roundworms, lungworm, eyeworm, sucking and biting lice, sarcoptic and chorioptic mange mites, buffalo fly and cattle tick in cattle, and for the treatment and control of gastrointestinal roundworms and lungworm of red deer.

PRODUCT BENEFITS*

- ✓ Controls worms, lice, cattle tick and buffalo fly;
- ✓ Convenience of a Pour-On application;
- ✓ Wide margin of safety when used as directed.
- ✓ Rainfast;
- ✓ Buffalo fly control - up to 21 days;
- ✓ Longest cattle tick control - 28 days[†];
- ✓ Longest lice protection - up to 56 days[†];

BROAD-SPECTRUM PARASITE CONTROL

When used as directed IVOMEC Pour-On treats and controls the following parasites of cattle:

Gastrointestinal roundworms:

Adult and immature: (L ₄):	Adult stages of:	Immature stages of:
Small brown stomach worm (including inhibited stages)	<i>Ostertagia ostertagi</i>	<i>Cooperia oncophora</i> <i>C. punctata</i>
Barbers pole worm	<i>Haemonchus placei</i>	<i>Oesophagostomum venulosum</i>
Stomach hair worm	<i>Trichostrongylus axei</i>	<i>Trichuris spp.</i>
Intestinal hair worm	<i>T. colubriformis</i>	<i>Strongyloides papillosum</i>
Small intestinal worm	<i>Cooperia spp.</i>	
Nodule worm	<i>Oesophagostomum radiatum</i>	

Lungworm: *Dictyocaulus viviparous* (adult and immature)

Eyeworm: *Thelazia* spp (adult stages)

Sucking and Biting Lice: *Linognathus vituli*
Haematopinus eurysternus
Bovicola (Damalinia) bovis
Solenopotes capillatus

Buffalo Fly: *Haematobia irritans exigua*

Mites: *Chorioptes bovis*

Sarcoptes scabiei var. *bovis*

Cattle Tick: *Boophilus microplus* (including organophosphate-, synthetic pyrethroid-, and amidine-resistant strains.)



PERSISTENT ACTIVITY

At the recommended dose of 1mL per 10kg, IVOMEC Pour-On controls:

- ✓ *Trichostrongylus axei* and *Cooperia* spp. for up to 14 days after treatment;
- ✓ *Ostertagia* spp. and *Oesophagostomum radiatum* for up to 21 days after treatment;
- ✓ *Dictyocaulus viviparous* (Lungworm) for up to 28 days after treatment;
- ✓ *Haematobia irritans exigua* (Buffalo Fly) for up to 21 days after treatment;
- ✓ Lice (*Bovicola bovis* and *Linognathus vituli*) for up to 56 days after treatment;
- ✓ Prevents the appearance of viable adult cattle ticks (*Boophilus microplus*) on cattle for 28 days after treatment.

DOSE RATE

1mL for each 10kg liveweight.

PACK SIZES

Available in 1 Litre, 2.5 Litre, 5 Litre & 20 Litre packs.

The 20L pack size is packaged in a robust drum for the larger producer.

Both the 2.5L and 5L pack sizes are packaged in convenient, collapsible backpacks for easy use.

The 1L pack size features the flexible 'Magicap' dispenser which eliminates the need for a drench gun.

Live Weight (kg)	Dose Volume (mL)	Dose			
		1L	2.5L	5L	20L
50-100	10	100	250	500	2000
101-150	15	150	375	750	3000
151-200	20	200	500	1000	4000
201-250	25	250	625	1250	5000
251-300	30	300	750	1500	6000
301-350	35	350	875	1750	7000
351-400	40	400	1000	2000	8000
401-450	45	450	1125	2250	9000
451-600	50	500	1250	2500	10000
501-650	55	550	1375	2750	11000
551-600	60	600	1500	3000	12000
601-650	65	650	1625	3250	13000

Heavier animals (eg mature bulls) should receive an additional 5mL for each 50kg over 650kg.

WITHHOLDING PERIOD

Meat: DO NOT APPLY to cattle less than 42 days before slaughter for human consumption.

DO NOT APPLY to deer less than 35 days before slaughter for human consumption.

Milk: When milk or milk products are to be used for human consumption do not administer to cows within 28 days prior to calving or during lactation.

ESI (Export Slaughter Interval)^Δ: 42 days

STORAGE

Store below 30°C (Room Temperature).

Store bottle in carton to protect from light.

*See product labels for full claim details. [†]Of the endectocide pour-on cattle products.

^ΔThe ESI was correct at the time of publication. Before using this product, confirm the current ESI from the APVMA website (www.apvma.gov.au/residues/ESI.shtml) or call Merial: 1800-808-691.

Merial Australia Pty Ltd, Building D, 12-24 Talavera Rd, Macquarie Park, NSW 2113 (ABN 53 071 187 285). [®]IVOMEC and the cattle head are registered trademarks of Merial Limited. ©2012 Merial Limited. All rights reserved. IVPO-12-000. Adze 5015

