IODINE WOUND SPRAY

By AgriLabs

Topical Wound Dressing

Indications: A topical antiseptic for use on horses, cattle, swine and sheep prior to surgical procedures such as castrating and docking, for application to the navel of newborn animals; and as an aid in the treatment of minor cuts, bruises and abrasions.

Dosage and Administration: Hold the container approximately four (4) to six (6) inches from the area to be treated. Point the valve at the area to be sprayed. Pull the trigger of the valve and spray the area once lightly. May be repeated once a day, when necessary, until abraded area is healed.

Note: When sprayed on or near the teats or udders of dairy animals, the teats and udders should be thoroughly washed before the next milking to prevent contamination of milk.

Contraindication(s): Not for use on burns or in body cavities or in deep wounds.

Precaution(s): Store in a cool place. Flammable. Do not expose to heat or store at a temperature above 120°F. Do not use near an open flame.

Caution(s): If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian. Harmful if swallowed. Do not apply to the eyes, mucous membranes or large areas of abraded skin. Avoid inhalation of mist.

Livestock remedy. Not for human use. Keep out of the reach of children.

Presentation: 16 fl. oz. (473 mL) and 1 gallon (3.785 L) bottles.